

Effects of weapon types, interventions, and transport times on complications in combat injuries to musculoskeletal system

 Rahman Senocak,¹  Huseyin Tas,²  Orhan Ureyen,³  Sahin Kaymak,¹  Oguz Hancerliogullari¹

¹Department of General Surgery, Gulhane Training and Research Hospital, Ankara, Turkey

²Department of General Surgery, Izmir Katip Celebi Universitesi Ataturk Training and Research Hospital, Izmir, Turkey

³Department of General Surgery, Bozyaka Training and Research Hospital, Izmir, Turkey

ABSTRACT

OBJECTIVE: To analyze the effects of weapon type, medical interventions, and transportation time on complications due to combat-related injuries of the musculoskeletal and soft tissue (ST).

METHODS: A total of 108 patients treated at the emergency department (ED) over a period of 3.5 years were included in this study. The effects of weapon type, type of first intervention team, interventions at the ED, and transport time on complications were compared by retrospectively assessing patients' recorded data.

RESULTS: The average age of 108 patients with ST injuries was 24 years. 25 patients developed complications. The ratio of complications in patients with injuries from explosive weapons was 25.42% and was significantly greater ($p<0.05$) than the rate observed from non-explosive weapons. Regarding the transport time of patients reaching the hospital, the greatest complication rate was 30.77% in patients transported to the hospital in 2–4 h; this rate was significantly ($p<0.05$) greater than for those transported in other time ranges. Regarding the ED procedures, the complication rate was 37.50% and significantly higher ($p<0.05$) in the group that received debridement + irrigation + late primary repair.

CONCLUSION: Transporting the injured to the ED within the first 2 h, treatment including irrigation and debridement, and secondary healing following explosive injuries containing contamination and dead tissue appear to be good options for reducing complications.

Keywords: Combat-related injuries; complication rates; musculoskeletal system; soft tissue.

Cite this article as: Senocak R, Tas H, Ureyen O, Kaymak S, Hancerliogullari O. Effects of weapon types, interventions, and transport times on complications in combat injuries to musculoskeletal system. *North Clin Istanbul* 2019;6(1):64–68.

Combat injuries of the musculoskeletal system (MSS) in today's battles are as high as 60–70% [1]. Approximately 40% of these injuries occur from bullets and 60% occur from the blast effect and fragmentation of explosive weapons [2, 3]. Explosive weapons are preferred by terrorist groups due to their ease of production, they are relatively inexpensive, and they are able to inflict numerous injuries and death.

In addition to the direct effects of explosive weapons,

injuries of MSS also occur with blast effects, in which broken pieces of surrounding objects are dragged into soft and bony tissues. From the beginning, this type of injury is typically more contaminated than projectile injuries, thus, increasing the complication rates and sequelae [2-5]. The effectiveness of wound debridement and irrigation, antibiotherapy, and surgical procedures on complication rates is well known [6-9]. However, there are some other factors affecting the complications

Received: August 23, 2017 *Accepted:* January 21, 2018 *Online:* August 07, 2018

Correspondence: Dr. Sahin KAYMAK, Saglik Bilimleri Universitesi, Gulhane Egitim ve Arastirma Hastanesi, Genel Cerrahi Klinigi, Ankara, Turkey.

Tel: +90 312 304 20 00 e-mail: sahin Kaymak@hotmail.com

© Copyright 2019 by Istanbul Provincial Directorate of Health - Available online at www.northclinist.com



of MSSs related to projectile injuries. According to various studies, it is still controversial whether the type of weapon, the person who makes the first intervention, the transport time, and the types of intervention at the emergency department (ED) have important effects on complications [8, 10].

The aim of this study was to present the effects of weapon type, type of first intervention team, interventions at the ED, and transport time on complications due to combat-related injuries of musculoskeletal and soft tissue (ST), which is the most common injury type in modern battles.

MATERIALS AND METHODS

Included in this study were 108 patients with combat injuries of the ST and MSS who were treated at Role II military hospital over a period of 3.5 years. A retrospective review of hospital records for all patients who sustained injuries was approved by the institutional review board. Patients with fractures, major vascular injuries, and penetrating injuries of the head, chest, and abdomen were excluded from the study.

The first interventions made during the pre-hospital period were carried out by paramedics when there was an emergency medical service available. Under other conditions, interventions were carried out by non-paramedics including those who were well-trained in the first aid. Patient transportation by air or highway was decided in accordance with the weather conditions and the general condition of the patient after communication with the ED.

Quick and effective systemic examinations were performed on the patients brought to the ED. At first, vital signs were evaluated, then vascular access was established, and blood sampling was made according to laboratory investigations that included a blood group determination. Anal region examinations and the urethral catheterizations of patients were also performed in patients with abdominal injuries. Prophylactic tetanus vaccines and intravenous isotonic solution with triple antibiotic therapy (first-generation cephalosporins, aminoglycosides, and metronidazole) were then given to all cases, and the patients were subsequently monitored. Because we are a Role 2 hospital, during the blood transfusion given to the cases at the shock table, whole blood was used after cross-matches per patient.

Typing wounds were made according to the injured tissues Type ST (wounds), Type F (wounds with fractures), Type V (vital wounds putting the patient's life

at risk), and Type VF (wounds with fractures and involving vital structures putting life or limb at risk). The wound was graded according to severity by the International Committee of the Red Cross. Grade 1 (Low energy transfer), Grade 2 (High energy transfer), Grade 3 (Massive energy transfer).

Superficial ST wounds required only simple local wound toilet. They were cleansed with soap, water, and a disinfectant and a simple dry dressing applied; the small wound was left open to heal by secondary intention. Grade 2 and 3 wounds required a full surgical exploration and excision, some required serial debridement or leave open, and late primary repair except ST of hand and head.

If <4 h since wounding, immediate primary closure, after full debridement, with a subcutaneous drain, was permissible. If >4 h old, it was better to leave the wound open for delayed primary closure after 2–4 days. This was usually accomplished by direct suture: Simple approximation of the deep structures and skin with minimal mobilization of the skin edges and without tension. Once the wound was adequately excised, it was covered with a bulky absorbent dressing made of absorbent cotton wool.

Surgical interventions such as wound dressings, primary repairs, irrigation + debridement + late primary repair, and irrigation + debridement + secondary healing were applied to patients in stable conditions. In our ED, the weapon type causing the injury, the person who provided first aid, the transport time, and the treatments performed at the ED were recorded.

Patients' data were evaluated retrospectively, and the effects of the weapon type causing the injury, the type of first treatment team, the interventions made in the ED, and transport times on the complications were compared. The SPSS for Windows 15.0 program (SPSS, Inc., Chicago, IL, USA) was used for statistical analyses. For comparisons among groups, Student's t-tests and Mann–Whitney U-tests were applied. Relationships of numeric values were investigated with correlation analyses. Qualitative data were compared with Chi-square tests. In statistical decisions, $p < 0.05$ level was accepted as a statistically significant difference.

RESULTS

The average age of the 108 patients evaluated in this study was 24 years. There were 91 extremity muscu-

loskeletal injuries, three abdominal injuries, two chest injuries, and 12 head injuries (Table 1).

Among the patients included in the study, complications developed in 25 patients, 20 (80%) of whom had wound infections. Other complications included one patient with a colostomy construction for protecting the wound, two had a loss of extremity function, one had a secondarily applied amputation due to compartment syndrome and wound infection, and one had orchitis (Table 2).

Comparisons among complications, type of weapon causing the injury, interventions in the ED, and transport times are shown in Table 3. According to these data, 54.6% of injuries were due to explosive weapons and 45.4% of injuries were due to bullets. However, complication rates were 25.42% for injuries with explosive weapons and only 20.41% in bullet-related injuries ($p < 0.05$). In comparing complication rates in terms of whether the first intervention was made by paramedics or not, there was no significant difference ($p > 0.05$). With regard to arrival times, 42.6% of patients arrived at the hospital within the 1st h, 38.9% arrived in 1–2 h, 12% in 2–4 h, and 6.5% in 4 h. There was a significantly higher complication rate (30.77%) for those patients who arrived within the 2–4 h period ($p < 0.05$). Treatment procedures in the ED included wound dressing to 37.04% of patients, primary repair to 6.48% of patients, irrigation + debridement + late primary repair to 22.22% of patients, and irrigation + debridement + secondary healing to 34.26% of patients. A significantly higher complication rate (37.5%) was observed in the irrigation + debridement + late primary repair group ($p < 0.05$).

DISCUSSION

The purpose of weapons in modern warfare is to inflict damage on the military battlefield. At present, the rate of explosive weapons usage is greater than the rate of bullet usage (60% vs. 40%, respectively) [2]. Likewise, explosive weapons are mostly preferred by terrorist groups because they can quickly cause numerous injuries and deaths. According to the study in which we have reported results from conflict zones, 54.6% of injuries were related to explosive weapons and 45.4% were related to bullets. This suggests that explosive weapons, whether they are used in a battlefield zone or terrorist zone, are much preferred in modern wars.

Many studies have examined complications with MSS

TABLE 1. Demographic data of the patients and anatomic locations of MSS-ST injuries

	Average	n	%
Age (years)	24		
Male		102	94.4
Female		6	5.6
Anatomic locations of MSS-ST GSW			
Abdomen		3	2.78
Thorax		2	1.85
Head		12	11.11
Only MSS-ST injuries		91	84.26

MSS: Musculoskeletal system; ST: Soft tissue.

TABLE 2. Types and rates of various complications

	Quantity	%
Wound infection	20	80
Diverting colostomy	1	4
Extremity dysfunction	2	8
Amputation [†]	1	4
Orchitis [‡]	1	4
Total	25	100

[†]Secondary to wound infection and compartment syndrome; [‡]Secondary to wound infection.

injuries, which comprise 60–70% of injuries occurring in today's battles where the usage of explosive weapons has increased [9–11]. These studies also indicate that complication rates are much higher for MSS injuries inflicted with high kinetic energy weapons [11]. In our study, the complication rate was 23.5%, and all patients were injured by high kinetic energy bullets or mortar shells or high kinetic energy weapons like antitank mines. Explosive weapons, due to their high levels of kinetic energy and blast effect, produce musculoskeletal injuries that tend to become contaminated. Therefore, complication rates are expected to be much higher (25.42%) than with bullet injuries.

In studies related to who provides the first intervention to trauma patients, complication rates have been shown to be less frequent in patients who first receive paramedic intervention [12]. In our study, although there were no sta-

TABLE 3. Comparisons among weapon types, first aid team, transport time to the hospital, and ED procedures

	Number of patients		Complication	
	n	%	n	%
Type of weapon				
Bullet	49	45.4	10	20.41
Explosive	59	54.6	15	25.42*
First aid team				
Paramedic	61	56.4	15	24.59
Non-paramedic	47	43.6	10	21.28
Transport time to hospital (hr)				
0–1	46	42.6	10	21.74
1–2	42	38.9	10	23.80
2–4	13	12.0	4	30.77*
>4	7	6.5	1	14.29
Procedure applied at the ED				
Surgical dressing	40	37.04	5	12.50
Primary suture	7	6.48	1	14.29
Irrigation + debridement + late primary suture	24	22.22	9	37.50*
Irrigation + debridement + secondary wound healing	37	34.26	10	27.03

*Significantly different than other groups ($p < 0.05$).

tistically significant differences between these two groups, complication rates were less frequent for interventions made by non-paramedics. However, we think this may be due to several reasons. For instance, this reduced rate may be due to limited interventions in battlefield zones, better trained first aid teams, and most importantly, sending the patient to the nearest health-care team by aircraft for those who are heavy trauma cases.

It is well known that complication rates increase when patient transportation times to the ED are prolonged [13, 14]. Moreover, some studies have reported that complication rates are increased if late debridement is delayed [10]. In our study, we generally found that complication rates increased as transportation times increased. A 30.77% complication rate was found in patients who were brought back within 2–4 h. Although slightly wounded patients who did not require irrigation and debridement reached the ED in times longer than 4 h due to transport by highway, a low complication rate of 14.29% was detected.

When surgical procedures applied at the ED were investigated for this study, only wound dressing or primary repair was applied to musculoskeletal injuries not

requiring debridement, and low complication rates were found. Irrigation + debridement + late primary repair or irrigation + debridement + secondary healing procedures were applied to ST and musculoskeletal injuries including wide, contaminated, highly dirty, and devitalized tissue. In many studies [6] containing these types of musculoskeletal injuries, if the skin is covered without compartment syndrome, the irrigation + debridement + late primary repair method is suggested. Under these circumstances, we observed a complication rate of 37.50%, which is statistically significant. We conclude that this significant difference arises from the fact that all patients in this study were exclusively injured by high kinetic energy weapons.

Conclusion

The first intervention made by paramedics or non-paramedics, but well-trained personnel does not create a significant difference in terms of complication development in conflict zones where medical intervention is limited. Bringing wounded patients to the ED within 2 hours and performing irrigation + debridement + secondary healing are a good option for reducing the com-

plications resulting from injuries from high kinetic energy weapons.

Acknowledgement: All authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Conflict of Interest: The authors declare no conflict of interest.

Financial Disclosure: The authors declared that this study has received no financial support.

Authorship contributions: Concept – H.T., O.U.; Design – H.T., O.U.; Data collection &/or processing – H.T., O.U.; Analysis and/or interpretation – H.T., O.U.; Writing – H.T., S.K., R.S. O.H.; Critical review – H.T., S.K., R.S.

REFERENCES

1. Covey DC. Blast and fragment injuries of the musculoskeletal system. *J Bone Joint Surg Am* 2002;84-A:1221–34. [CrossRef]
2. Ritenour AE, Blackburn LH, Kelly JF, McLaughlin DF, Pearse LA, Holcomb JB, et al. Incidence of primary blast injury in US military overseas contingency operations: A retrospective study. *Ann Surg* 2010;251:1140–4. [CrossRef]
3. Özpek A, Hasbahçeci M, Kabak İ, Yücel M, Çalıřkan M, Alimođlu O. Therapeutic approaches to the abdominal gunshot wounds. *Turk J Surg* 2012;28:76–80. [CrossRef]
4. Hill PF, Edwards DP, Bowyer GW. Small fragment wounds: Bio-physics, pathophysiology and principles of management. *J R Army Med Corps* 2001;147:41–51. [CrossRef]
5. Bartlett CS, Helfet DL, Hausman MR, Strauss E. Ballistics and gunshot wounds: Effects on musculoskeletal tissues. *J Am Acad Orthop Surg* 2000;8:21–36. [CrossRef]
6. Mérens A, Rapp C, Delaune D, Danis J, Berger F, Michel R, et al. Prevention of combat-related infections: Antimicrobial therapy in battlefield and barrier measures in French military medical treatment facilities. *Travel Med Infect Dis* 2014;12:318–29. [CrossRef]
7. Murray CK, Obremskey WT, Hsu JR, Andersen RC, Calhoun JH, Clasper JC, et al. Prevention of infections associated with combat-related extremity injuries. *J Trauma* 2011;71:S235–57. [CrossRef]
8. Manning MM, Hawk A, Calhoun JH, Andersen RC. Treatment of war wounds: A historical review. *Clin Orthop Relat Res* 2009;467:2168–91.
9. Murray CK, Hsu JR, Solomkin JS, Keeling JJ, Andersen RC, Ficke JR, et al. Prevention and management of infections associated with combat-related extremity injuries. *J Trauma* 2008;64:S239–51. [CrossRef]
10. Tař H, Mesci A, Eryılmaz M, Zeybek N, Peker Y. The affecting factors on the complication ratio in abdominal gunshot wounds. *Ulus Travma Acil Cerrahi Derg* 2011;17:450–4. [CrossRef]
11. Persad IJ, Reddy RS, Saunders MA, Patel J. Gunshot injuries to the extremities: Experience of a U.K. Trauma centre. *Injury* 2005;36:407–11.
12. Murad MK, Larsen S, Husum H. Prehospital trauma care reduces mortality. Ten-year results from a time-cohort and trauma audit study in Iraq. *Scand J Trauma Resusc Emerg Med* 2012;20:13. [CrossRef]
13. Hebrang A, Henigsberg N, Hrabac P. Evacuation times of civilians and soldiers wounded during the war in Croatia. *Mil Med* 2006;171:1045–50.
14. Leppäniemi AK. Abdominal war wounds-experiences from red cross field hospitals. *World J Surg* 2005;29 Suppl 1:S67–71. [CrossRef]